



Boundaries of Levi-flat hypersurfaces: special hyperbolic points

Pierre Dolbeault

► To cite this version:

Pierre Dolbeault. Boundaries of Levi-flat hypersurfaces: special hyperbolic points. *Annales Polonici Mathematici*, 2012, 106, pp.145-170. 10.4064/ap106-0-12 . hal-00669176v2

HAL Id: hal-00669176

<https://hal.science/hal-00669176v2>

Submitted on 7 Jan 2013

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

BOUNDARIES OF LEVI-FLAT HYPERSURFACES: SPECIAL HYPERBOLIC POINTS

PIERRE DOLBEAULT

ABSTRACT. Let $S \subset \mathbb{C}^n$, $n \geq 3$ be a compact connected 2-codimensional submanifold having the following property: there exists a Levi-flat hypersurface whose boundary is S , possibly as a current. Our goal is to get examples of such S containing at least one special 1-hyperbolic point: sphere with two horns; elementary models and their gluing. The particular cases of graphs are also described.

1. INTRODUCTION

Let $S \subset \mathbb{C}^n$, be a compact connected 2-codimensional submanifold having the following property: there exists a Levi-flat hypersurface $M \subset \mathbb{C}^n \setminus S$ such that $dM = S$ (i.e. whose boundary is S , possibly as a current). The case $n = 2$ has been intensively studied since the beginning of the eighties, in particular by Bedford, Gaveau, Klingenberg; Shcherbina, Chirka, G. Tomassini, Slodkowski, Gromov, Eliashberg; it needs global conditions: S has to be contained in the boundary of a strictly pseudoconvex domain.

We consider the case $n \geq 3$; results on this case has been obtained since 2005 by Dolbeault, Tomassini and Zaitsev, local necessary conditions recalled in section 2 have to be satisfied by S , the singular CR points on S are supposed to be elliptic and the solution M is obtained in the sense of currents [DTZ05, DTZ10]. More recently a regular solution M has been obtained when S satisfies a supplementary global condition as in the case $n = 2$ [DTZ09], the singular CR points on S still supposed to be elliptic.

The problem we are interested in is to get examples of such S containing at least one special 1-hyperbolic point (section 2.4). The CR-orbits near a special 1-hyperbolic point are large and, assuming them compact, a careful examination has to be done (sections 2.6, 2.7). As a topological preliminary, we need a generalization of a theorem of Bishop on the difference of the numbers of special elliptic and 1-hyperbolic points (section 2.8); this result is a particular case of a theorem of Hon-Fei Lai [Lai72].

The first considered example is the sphere with two horns which has one special 1-hyperbolic point and three special elliptic points (section 3.4). Then we consider elementary models and their gluing to obtain more complicated examples (section 3.5). Results have been announced in [Dol08], and

in more precise way in [Dol11]; the first aim of this paper is to give complete proofs. Finally, we recall in detail and extend the results of [DTZ09] on regularity of the solution when S is a graph satisfying a supplementary global condition, as in the case $n = 2$, to the case of existence of special 1-hyperbolic points, and to gluing of elementary smooth models (section 4).

2. PRELIMINARIES: LOCAL AND GLOBAL PROPERTIES OF THE BOUNDARY

2.1. Definitions. A smooth, connected, CR submanifold $M \subset \mathbb{C}^n$ is called *minimal* at a point p if there does not exist a submanifold N of M of lower dimension through p such that $HN = HM|_N$. By a theorem of Sussman, all possible submanifolds N such that $HN = HM|_N$ contain, as germs at p , one of the minimal possible dimension, defining a so called CR *orbit* of p in M whose germ at p is uniquely determined.

Let S be a smooth compact connected oriented submanifold of dimension $2n - 2$. S is said to be a *locally flat boundary* at a point p if it locally bounds a Levi-flat hypersurface near p . Assume that S is CR in a small enough neighborhood U of $p \in S$. If all CR orbits of S are 1-codimensional (which will appear as a necessary condition for our problem), the following two conditions are equivalent [DTZ05]:

- (i) S is a locally flat boundary on U ;
- (ii) S is nowhere minimal on U .

2.2. Complex points of S . (i.e. singular CR points on S) [DTZ05].

At such a point $p \in S$, $T_p S$ is a complex hyperplane in $T_p \mathbb{C}^n$. In suitable local holomorphic coordinates $(z, w) \in \mathbb{C}^{n-1} \times \mathbb{C}$ vanishing at p , with $w = z_n$ and $z = (z_1, \dots, z_{n-1})$, S is locally given by the equation

$$(1) \quad w = Q(z) + O(|z|^3), \quad Q(z) = \sum_{1 \leq i, j \leq n-1} (a_{ij} z_i z_j + b_{ij} z_i \bar{z}_j + c_{ij} \bar{z}_i \bar{z}_j)$$

S is said *flat* at a complex point $p \in S$ if $\sum b_{ij} z_i \bar{z}_j \in \lambda \mathbf{R}$, $\lambda \in \mathbb{C}$. We also say that p is *flat*.

Let $S \subset \mathbb{C}^n$ be a locally flat boundary with a complex point p . Then p is *flat*.

By making the change of coordinates $(z, w) \mapsto (z, \lambda^{-1} w)$, we get $\sum b_{ij} z_i z_j \in \mathbf{R}$ for all z . By a change of coordinates $(z, w) \mapsto (z, w + \sum a'_{ij} z_i z_j)$ we can choose the holomorphic term in (1) to be the conjugate of the antiholomorphic one and so make the whole form Q real-valued.

We say that S is in a *flat normal form* at p if the coordinates (z, w) as in (1) are chosen such that $Q(z) \in \mathbf{R}$ for all $z \in \mathbb{C}^{n-1}$.

2.2.1. Properties of Q . Assume that S is in a flat normal form; then, the quadratic form Q is real valued. If Q is positive definite or negative definite, the point $p \in S$ is said to be *elliptic*; if the point $p \in S$ is not elliptic, and if Q is non degenerate, p is said to be *hyperbolic*. From section 2.4, we will only consider particular cases of the quadratic form Q .

2.3. Elliptic points.

2.3.1. Properties of Q .

Proposition 1. ([DTZ05, DTZ10]). *Assume that $S \subset \mathbb{C}^n$, ($n \geq 3$) is nowhere minimal at all its CR points and has an elliptic flat complex point p . Then there exists a neighborhood V of p such that $V \setminus \{p\}$ is foliated by compact real $(2n - 3)$ -dimensional CR orbits diffeomorphic to the sphere \mathbb{S}^{2n-3} and there exists a smooth function ν , having the CR orbits as the level surfaces.*

Sketch of Proof. (see [DTZ10]). In the case of a quadric S_0 ($w = Q(z)$), the CR orbits are defined by $w_0 = Q(z)$, where w_0 is constant. Using (1), we approximate the tangent space to S by the tangent space to S_0 at a point with the same coordinate z ; the same is done for the tangent spaces to the CR orbits on S and S_0 ; then we construct the global CR orbit on S through any given point close enough to p . \square

2.4. Special flat complex points. From [Bis65], for $n = 2$, in suitable local holomorphic coordinates centered at 0, $Q(z) = (z\bar{z} + \lambda \operatorname{Re} z^2)$, $\lambda \geq 0$, under the notations of [BK91]; for $0 \leq \lambda < 1$, p is said to be *elliptic*, and for $1 < \lambda$, it is said to be *hyperbolic*. The parabolic case $\lambda = 1$, not generic, will be omitted [BK91]. When $n \geq 3$, the Bishop's reduction cannot be generalized.

We say that the flat complex point $p \in S$ is *special* if in convenient holomorphic coordinates centered at 0,

$$(2) \quad Q(z) = \sum_{j=1}^{n-1} (z_j \bar{z}_j + \lambda_j \operatorname{Re} z_j^2), \quad \lambda_j \geq 0$$

Let $z_j = x_j + iy_j$, x_j, y_j real, $j = 1, \dots, n-1$, then:

$$(3) \quad Q(z) = \sum_{l=1}^{n-1} ((1 + \lambda_l)x_l^2 + (1 - \lambda_l)y_l^2) + O(|z|^3).$$

A flat point $p \in S$ is said to be *special elliptic* if $0 \leq \lambda_j < 1$ for any j .

A flat point $p \in S$ is said to be *special k -hyperbolic* if $1 < \lambda_j$ for $j \in J \subset \{1, \dots, n-1\}$ and $0 \leq \lambda_j < 1$ for $j \in \{1, \dots, n-1\} \setminus J \neq \emptyset$, where k denotes the number of elements of J .

Special elliptic (resp. special k -hyperbolic) points are elliptic (resp. hyperbolic).

Special flat complex points

2.5. Special hyperbolic points. S being given by (1), let S_0 be the quadric of equation $w = Q(z)$.

Lemma 2. *Suppose that S_0 is flat at 0 and that 0 is a special k -hyperbolic point. Then, in a neighborhood of 0, and with the above local coordinates, S_0 is CR and nowhere minimal outside 0, and the CR orbits of S_0 are the $(2n - 3)$ -dimensional submanifolds given by $w = \operatorname{const.} \neq 0$.*

Proof. The submanifolds $w = \text{const.} \neq 0$ have the same complex tangent space as S_0 and are of minimal dimension among submanifolds having this property, so they are CR orbits of codimension 1, and from the end of section 2.1, S_0 is nowhere minimal outside 0.

The section $w = 0$ of S_0 is a real quadratic cone Σ'_0 in \mathbf{R}^{2n} whose vertex is 0 and, outside 0, it is a CR orbit Σ_0 in the neighborhood of 0. We will improperly call Σ'_0 a *singular CR orbit*. \square

2.6. Foliation by CR-orbits in the neighborhood of a special 1-hyperbolic point. We first mimic and transpose the beginning of the proof of Proposition 1, i.e. of 2.4.2. in ([DTZ05, DTZ09]).

2.6.1. *Local 2-codimensional submanifolds.* In order to use simple notations, we will assume $n = 3$.

In \mathbb{C}^3 , consider the 4-dimensional submanifold S locally defined by the equation

$$(1) \quad w = \varphi(z) = Q(z) + O(|z|^3)$$

and the 4-dimensional submanifold S_0 of equation

$$(4) \quad w = Q(z)$$

with

$$Q = (\lambda_1 + 1)x_1^2 - (\lambda_1 - 1)y_1^2 + (1 + \lambda_2)x_2^2 + (1 - \lambda_2)y_2^2$$

having a special 1-hyperbolic point at 0, ($\lambda_1 > 1, 0 \leq \lambda_2 < 1$), and the cone Σ'_0 whose equation is: $Q = 0$. On S_0 , a CR orbit is the 3-dimensional submanifold \mathcal{K}_{w_0} whose equation is $w_0 = Q(z)$. If $w_0 > 0$, \mathcal{K}_{w_0} does not cut the line $L = \{x_1 = x_2 = y_2 = 0\}$; if $w_0 < 0$, \mathcal{K}_{w_0} cuts L at two points.

Lemma 3. $\Sigma_0 = \Sigma'_0 \setminus 0$ has two connected components in a neighborhood of 0.

Proof. The equation of $\Sigma'_0 \cap \{y_1 = 0\}$ is

$(\lambda_1 + 1)x_1^2 + (1 + \lambda_2)x_2^2 + (1 - \lambda_2)y_2^2 = 0$ whose only zero, in the neighborhood of 0, is $\{0\}$: the connected components are obtained for $y_1 > 0$ and $y_1 < 0$ respectively. \square

Local 2-codimensional submanifolds

2.6.2. *CR-orbits.* By differentiating (1), we get for the tangent spaces the following asymptotics

$$(5) \quad T_{(z, \varphi(z))}S = T_{(z, Q(z))}S_0 + O(|z|^2), \quad z \in \mathbb{C}^2$$

Here both $T_{(z, \varphi(z))}S$ and $T_{(z, Q(z))}S_0$ depend continuously on z near the origin.

Consider

(i) the hyperboloïd $H_- = \{Q = -1\}$, (then $Q(\frac{z}{(-Q(z))^{1/2}}) = -1$), and the projection:

$$\pi_- : \mathbb{C}^3 \setminus \{z = 0\} \rightarrow H_-, \quad (z, w) \mapsto \frac{z}{(-Q(z))^{1/2}},$$

(ii) for every $z \in H_-$, a real orthonormal basis $e_1(z), \dots, e_6(z)$ of $\mathbb{C}^3 \cong \mathbb{R}^6$ such that

$$e_1(z), e_2(z) \in H_z H_-, \quad e_3(z) \in T_z H_-,$$

where HH_- is the complex tangent bundle to H_- .

Locally such a basis can be chosen continuously depending on z . For every $(z, w) \in \mathbb{C}^3 \setminus \{z = 0\}$, consider the basis $e_1(\pi_-(z, w)), \dots, e_6(\pi_-(z, w))$. The unit vectors $e_1(\pi_-(z, w_0)), e_2(\pi_-(z, w_0)), e_3(\pi_-(z, w_0))$ are tangent to the CR orbit \mathcal{K}_{w_0} in (z, w_0) for $w_0 < 0$. Then, from (5), we have:

$$(6) \quad H_{(z, \varphi(z))} S = H_{(z, Q(z))} S_0 + O(|z|^2), \quad z \neq 0, \quad z \rightarrow 0.$$

As in [DTZ10], in the neighborhood of 0, denote by $E(q), q \in S \setminus \{0\}, w < 0$ the tangent space to the local CR orbit \mathcal{K} on S through q , and by $E_0(q_0), q_0 \in S_0 \setminus \{0\}, w < 0$ the analogous object for S_0 . We have :

$$(7) \quad E(z, \varphi(z)) = E_0(z, Q(z)) + O(|z|^2), \quad z \neq 0, \quad z \rightarrow 0$$

Given $\underline{q} \in S$, by integration of $E(q), q \in S$, we get, locally, the CR orbit (the leaf), on S through \underline{q} ; given $\underline{q}_0 \in S_0$, by integration of $E_0(q_0), q_0 \in S_0$, we get, locally, the CR orbit (the leaf), on S_0 through \underline{q}_0 (theorem of Sussman). On S_0 , a leaf is the 3-dimensional submanifold $\mathcal{K}_{\underline{q}_0} = \mathcal{K}_{w_0} = \mathcal{K}_0$ whose equation is $w_0 = Q(z)$, with $\underline{q} = (z_0, w_0 = Q(z_0))$. $d\pi_-$ projects each $E_0(q), q \in S_0, w < 0$, bijectively onto $T_{\pi(q)} H_-$, then $\pi_-|_{\mathcal{K}_0}$ is a diffeomorphism onto H_- ; this implies, from (7), that, in a suitable neighborhood of the origin, the restriction of π_- to each local CR orbit of S is a local diffeomorphism.

We have: $\varphi(z) = Q(z) + \Phi(z)$ with $\Phi(z) = O(|z|^3)$.

2.6.3. Behaviour of local CR orbits. Follow the construction of $E(z, \varphi(z))$; compare with $E_0(z, Q(z))$. We know the integral manifold, the orbit of $E_0(z, Q(z))$; deduce an evaluation of the integral manifold \mathcal{K} of $E(z, \varphi(z))$.

Lemma 4. *Under the above hypotheses, the local orbit Σ corresponding to Σ_0 has two connected components in the neighborhood of 0.*

Proof. Using the real coordinates, as for Lemma 3, $\Sigma' \cap \{y_1 = 0\}$. Locally, the connected components are obtained for $y_1 > 0$ and $y_1 < 0$ respectively, from formula (1). \square

We will improperly call $\Sigma' = \overline{\Sigma}$ a *singular CR orbit* and a *singular leaf of the foliation*.

We intend to prove: 1) \mathcal{K} does not cross the singular leaf through 0;

2) the only separatrix is the singular leaf through 0.

From the orbit \mathcal{K}_0 , construct the differential equation defining it, and using (7), construct the differential equation defining \mathcal{K} .

In \mathbb{C}^3 , we use the notations: $x = x_1, y = y_1, u = x_2, v = y_2$; it suffices to consider the particular case: $Q = 3x^2 - y^2 + u^2 + v^2$. On S_0 , the orbit \mathcal{K}_0 issued from the point $(c, 0, 0, 0)$ is defined by: $3x^2 - y^2 - u^2 + v^2 = 3c^2$, i.e., for $x \geq 0$, $x = \frac{1}{\sqrt{3}}(y^2 - u^2 - v^2 + 3c^2)^{\frac{1}{2}} = A(y, u, v)$; the local coordinates on the orbit are (y, u, v) . \mathcal{K}_0 satisfies the differential equation: $dx = dA$. From (9), the orbit \mathcal{K} , issued from $(c, 0, 0, 0)$, satisfies $dx = dA + \Psi$ with $\Psi(y, u, v; c) = O(|z|^2)$; hence $\Psi = d\Phi$, then $x = A + \Phi$, with $\Phi = O(|z|^3)$. More explicitly, \mathcal{K} is defined by:

$$x = x_{\mathcal{K},c} = \frac{1}{\sqrt{3}}(y^2 - u^2 - v^2 + 3c^2)^{\frac{1}{2}} + \Phi(y, u, v; c), \quad \Phi(y, u, v; c) = O(|z|^3)$$

The cone Σ'_0 whose equation is: $Q = 0$ is a separatrix for the orbits \mathcal{K}_0 . The corresponding object $\Sigma' = \{\varphi(z) = 0\}$ for S has the singular point 0 and for $x > 0, y > 0, u > 0, v > 0$ is defined by the differential equation $dx = d(A + \Phi)$, with $c = 0$, i.e. the local equation of Σ' is

$$x = x_{\mathcal{K},0} = \frac{1}{\sqrt{3}}(y^2 - u^2 - v^2)^{\frac{1}{2}} + \Phi(y, u, v; 0), \quad \Phi(y, u, v; 0) = O(|z|^3)$$

For given (y, u, v) , $x_{\mathcal{K},c} - x_{\mathcal{K},0} = x_{\mathcal{K}_0,c} - x_{\mathcal{K}_0,0} + \Phi(y, u, v; c) - \Phi(y, u, v; 0)$. But $x_{\mathcal{K}_0,c} - x_{\mathcal{K}_0,0} = O(1)$ and $\Phi(y, u, v; c) - \Phi(y, u, v; 0) = O(|z|^3)$.

As a consequence, for $x > 0, y > 0, u > 0, v > 0$, locally, Σ' is a separatrix for the orbits \mathcal{K} , and the only one. Same result for $x < 0$.

2.6.4. What has been done from the hyperboloid $H_- = \{Q = -1\}$ can be repeated from the hyperboloid $H_+ = \{Q = 1\}$.

As at the beginning of the section 2.6.2, we consider

(i) the hyperboloid $H_+ \{Q = 1\}$ and the projection:

$$\pi_+ : \mathbb{C}^3 \setminus \{z = 0\} \rightarrow H_+, \quad (z, w) \mapsto \frac{z}{(Q(z))^{1/2}},$$

(ii) for every $z \in H_+$, a real orthonormal basis $e_1(z), \dots, e_6(z)$ of $\mathbb{C}^3 \cong \mathbb{R}^6$ such that

$$e_1(z), e_2(z) \in H_z H_+, \quad e_3(z) \in T_z H_+,$$

where HH_+ is the complex tangent bundle to H_+ .

2.6.5.

Lemma 5. *Given φ , there exists $R > 0$ such that, in $B(0, R) \cap \{x > 0, y > 0, u > 0, v > 0\} \subset \mathbb{C}^2$, the CR orbits \mathcal{K} have Σ' as unique separatrix.*

Proof. When c tends to zero, $x_{\mathcal{K},c} - x_{\mathcal{K},0} = x_{\mathcal{K}_0,c} - x_{\mathcal{K}_0,0} = O(|z|)$, $\Phi(y, u, v; c) - \Phi(y, u, v; 0) = O(|z|^3)$. For $\varphi(z) = Q(z) + \Phi(z)$ with $\Phi(z) = O(|z|^3)$ given, in (9), $E(z, \varphi(z)) - E_0(z, Q(z)) = O(|z|^2)$ and $\Phi(y, u, v; c) - \Phi(y, u, v; 0) = O(|z|^3)$ are also given. Then there exists R such that, for $|z| < R$, $x_{\mathcal{K},c} - x_{\mathcal{K},0} > 0$. \square

2.7. CR-orbits near a subvariety containing a special 1-hyperbolic point.

2.7.1. In the section 2.7, we will impose conditions on S and give a local property in the neighborhood of a compact $(2n - 3)$ -subvariety of S .

Assume that $S \subset \mathbb{C}^n$ ($n \geq 3$), is a locally closed $(2n - 2)$ -submanifold, nowhere minimal at all its CR points, which has a unique 1-hyperbolic flat complex point p , and such that:

(i) Σ being the orbit whose closure Σ' contains p , then Σ' is compact.

Let $q \in S$, $q \neq p$; then, in a neighborhood U of q disjoint from p , S is CR, $\text{CR-dim } S = n - 2$, S is non minimal and Σ is 1-codimensional. To show that the CR orbits constitute a foliation on S whose separatrix is Σ' : this is true in U since $\Sigma \cap U$ is a leaf. Moreover, let U_0 the ball $B(0, R)$ centered in $p = 0$ in Lemma 5, if $U \cap U_0 \neq \emptyset$, the leaves in U glue with the leaves in U_0 on $U \cap U_0$. Since Σ' is compact, there exists a finite number of points $q_j \in \Sigma'$, $j = 0, 1, \dots, J$, and open neighborhoods U_j , as above, such that $(U_j)_{j=0}^J$ is an open covering of Σ' . Moreover the leaves on U_j glue respectively with the leaves on U_k if $U_j \cap U_k \neq \emptyset$.

2.7.2.

Proposition 6. *Assume that $S \subset \mathbb{C}^n$ ($n \geq 3$), is a locally closed $(2n - 2)$ -submanifold, nowhere minimal at all its CR points, which has a unique special 1-hyperbolic flat complex point p , and such that:*

(i) Σ being the orbit whose closure Σ' contains p , then Σ' is compact;

(ii) Σ has two connected components σ_1, σ_2 , whose closures are homeomorphic to spheres of dimension $2n - 3$.

Then, there exists a neighborhood V of Σ' such that $V \setminus \Sigma'$ is foliated by compact real $(2n - 3)$ -dimensional CR orbits whose equation, in a neighborhood of p is (3), and, the $w(= x_n)$ -axis being assumed to be vertical, each orbit is diffeomorphic to

the sphere \mathbf{S}^{2n-3} above Σ' ,

the union of two spheres \mathbf{S}^{2n-3} under Σ' ,

and there exists a smooth function ν , having the CR orbits as the level surfaces.

Proof. From subsection 2.7.1 and the following remark:

When x_n tends to 0, the orbits tends to Σ' , and because of the geometry of the orbits near p , they are diffeomorphic to a sphere above Σ' , and to the union of two spheres under Σ' . The existence of ν is proved as in Proposition 1, namely, consider a smooth curve $\gamma : [0, \varepsilon) \rightarrow S$ such that

$\gamma(0) = q$, where q is a point of Σ close to p , and γ is a diffeomorphism onto its image $\Gamma = \gamma([0, \varepsilon))$. Let $\nu = \gamma^{-1}$ on the image of γ , then, close enough to q , every CR orbit cuts Γ at a unique point $q(t)$, $t \in [0, \varepsilon)$. Hence there is a unique extension of ν from $\gamma([0, \varepsilon))$ to $V \setminus p$ where V is a neighborhood of Σ' having CR orbits as its level surfaces. ν being smooth away from p , it is smooth on the orbit Σ and, if we set $\nu(p) = \nu(q) = 0$, ν is smooth on a neighborhood of $\Sigma \cup \{p\} = \Sigma'$. \square

2.8. Geometry of the complex points of S . The results of section 2.8 are particular cases of theorems of H-F Lai [Lai72], that I learnt from F. Forstneric in July 2011.

In [BK91] E. Bedford & W. Klingenberg cite the following theorem of E. Bishop [Bis65][section 4, p.15]: *On a 2-sphere embedded in \mathbb{C}^2 , the difference between the numbers of elliptic points and of hyperbolic points is the Euler-Poincaré characteristic, i.e. 2.* For the proof, Bishop uses a theorem of ([CS 51], section 4).

We extend the result for $n \geq 3$ and give proofs which are essentially the same than in the general case of [Lai72, Lai74] but simpler.

2.8.1. Let S be a smooth compact connected oriented submanifold of dimension $2n - 2$. Let G be the manifold of the oriented real linear $(2n - 2)$ -subspaces of \mathbb{C}^n . The submanifold S of \mathbb{C}^n has a given orientation which defines an orientation $o(p)$ of the tangent space to S at any point $p \in S$. By mapping each point of S into its oriented tangent space, we get a smooth Gauss map

$$t : S \rightarrow G$$

Denote $-t(p)$ the tangent space to S at p with opposite orientation $-o(p)$.

2.8.2. Properties of G . (a) $\dim G = 2(2n - 2)$.

Proof. G is a two-fold covering of the Grassmannian $M_{m,k}$, of the linear k -subspaces of \mathbb{R}^m [Ste99][Part, section 7.9], for $m = 2n$ and $k = 2n - 2$; they have the same dimension. We have:

$$M_{m,k} \cong O_m/O_k \times O_{m-k}$$

But $\dim O_k = \frac{1}{2}k(k - 1)$, hence $\dim M_{m,k} = \frac{1}{2}(m(m - 1) - k(k - 1) - (m - k)(m - k - 1)) = k(m - k)$.

(b) G has the complex structure of a smooth quadric of complex dimension $(2n - 2)$ of $\mathbb{C}P^{2n-1}$ L74, [Pol08].

(c) There exists a canonical isomorphism $h : G \rightarrow \mathbb{C}P^{n-1} \times \mathbb{C}P^{n-1}$.

(d) Homology of G (cf [Pol08]): Let S_1, S_2 be generators of $H_{2n-2}(G, \mathbb{Z})$; we assume that S_1 and S_2 are fundamental cycles of complex projective subspaces of complex dimension $(n - 1)$ of the complex quadric G . We also denote S_1, S_2 the ordered two factors $\mathbb{C}P^{n-1}$, so that $h : G \rightarrow S_1 \times S_2$.

□

2.8.3.

Proposition 7. *For $n \geq 2$, in general, S has isolated complex points.*

Proof. Let $\pi \in G$ be a complex hyperplane of \mathbb{C}^n whose orientation is induced by its complex structure; the set of such π is $H = G_{n-1,n}^{\mathbb{C}} = \mathbb{CP}^{n-1*} \subset G$, as real submanifold. If p is a complex point of S , then $t(p) \in H$ or $-t(p) \in H$. The set of complex points of S is the inverse image by t of the intersections $t(S) \cap H$ and $-t(S) \cap H$ in G . Since $\dim t(S) = 2n - 2$, $\dim H = 2(n - 1)$, $\dim G = 2(2n - 2)$, the intersection is 0-dimensional, in general. □

2.8.4. Denoting also S , the fundamental cycle of the submanifold S and t_* the homomorphism defined by t , we have:

$$t_*(S) \sim u_1 S_1 + u_2 S_2$$

where \sim means *homologous to*.

2.8.5.

Lemma 8 (proved for $n = 2$ in [CS51]). *With the above notations, we have: $u_1 = u_2$; $u_1 + u_2 = \chi(S)$, Euler-Poincaré characteristic of S .*

The proof for $n = 2$ works for any $n \geq 3$, namely:

Let G' be the manifold of the oriented real linear 2-subspaces of \mathbb{C}^n . Let $\alpha : G \rightarrow G'$ map each oriented $2(n - 1)$ -subspace R onto its normal 2-subspace R' oriented so that R, R' determine the orientation of \mathbb{C}^n . α is a canonical isomorphism. Let $n : S \rightarrow G'$ the map defined by taking oriented normal planes; then: $n = \alpha t$ and $t = \alpha^{-1} n$, hence the mapping $h\alpha h^{-1} : S_1 \times S_2 \rightarrow S_1 \times S_2$. Let (x, y) be a point of $S_1 \times S_2$, then $(\dagger) \quad h\alpha h^{-1}(x, y) = (x, -y)$.

Over G , there is a bundle V of spheres obtained by considering as fiber over a real oriented linear $(2n - 2)$ -subspace of \mathbb{C}^n through 0 the unit sphere \mathbf{S}^{2n-3} of this subspace. Let Ω be the characteristic class of V , and let Ω_t, Ω_n denote the characteristic classes of the tangent and normal bundles of S . Then $t^*\Omega = \Omega_t, n^*\Omega = \Omega_n$.

V is the Stiefel manifold of ordered pairs of orthogonal unit vectors through in $\mathbb{R}^{2n} \cong \mathbb{C}^n$. Let $f : V \rightarrow G$ the projection.

From the Gysin sequence, we see that the kernel of $f^* : H^{2n-2}(G) \rightarrow H^{2n-2}(V)$ is generated by Ω . To find the kernel of f^* , we determine the morphism $f_* : H_{2n-2}(V) \rightarrow H_{2n-2}(G)$. A generating $2n - 2$ -cycle of in V is $S^2 \times e$ where $S^2 \cong \mathbb{CP}^{n-1}$ and e is a point. Let z be any point of S^2 , then from (\dagger) , we have

$$hf(z, e) = (z, -z)$$

Therefore, we see that $f_*(S^2 \times e) = S_1 - S_2$. Then, the kernel of f^* is \mathbb{Z} -generated by $S_1^* + S_2^*$.

With convenient orientation for the fibre of the bundle V , we get: $\Omega = S_1^* + S_2^*$. For convenient orientation of S , we get $\Omega_t.S = \chi_S = \text{Euler characteristic of } S$. We have

$$\Omega_t = t^*(S_1^* + S_2^*) = t^*S_1^* + t^*S_2^*$$

$$\Omega_n = n^*(S_1^* + S_2^*) = t^*\alpha^*(S_1^* + S_2^*) = t^*(S_1^* - S_2^*) = t^*S_1^* - t^*S_2^*$$

Since $\Omega_n = 0$, we get:

$$(t^*S_1^*).S = (t^*S_2^*).S = \frac{1}{2}\chi_S$$

2.8.6. Local intersection numbers of H and $t(S)$ when all complex points are flat and special. H is a complex linear $(n-1)$ -subspace of G , then is homologous to one of the S_j , $j = 1, 2$, say S_2 when G has its structure of complex quadric. The intersection number of H and S_1 is 1 and the intersection number of H and S_2 is 0. So, the intersection number of H and $u_1S_1 + u_2S_2$ is u_1 .

In the neighborhood of a complex point 0, S is defined by equation (1), with $w = z_n$ and

$$(1') \quad Q(z) = \sum_{j=1}^{n-1} \mu_j(z_j \bar{z}_j + \lambda_j \mathcal{R}e \ z_j^2), \quad \mu_j > 0, \lambda_j \geq 0$$

Let $z_j = x_{2j-1} + ix_{2j}$, $j = 1, \dots, n$, with real x_l . Let e_l the unit vector of the x_l axis, $l = 1, \dots, 2n$.

For simplicity assume $n = 3$: $Q(z) = \mu_1(z_1 \bar{z}_1 + \lambda_1 \mathcal{R}e \ z_1^2) + \mu_2(z_2 \bar{z}_2 + \lambda_2 \mathcal{R}e \ z_2^2)$, with $\mu_1 = \mu_2 = 1$.

Then, up to higher order terms, S is defined by:

$$z_1 = x_1 + ix_2; \quad z_2 = x_3 + ix_4; \quad z_3 = (1 + \lambda_1)x_1^2 + (1 - \lambda_1)x_2^2 + (1 + \lambda_2)x_3^2 + (1 - \lambda_2)x_4^2.$$

In the neighborhood of 0, the tangent space to S is defined by the four independent vectors

$$\begin{aligned} \nu_1 &= e_1 + 2(1 + \lambda_1)x_1 e_5; \quad \nu_2 = e_2 + 2(1 - \lambda_1)x_2 e_5; \quad \nu_3 = e_3 + 2(1 + \lambda_2)x_3 e_5; \\ \nu_4 &= e_4 + 2(1 - \lambda_2)x_4 e_5 \end{aligned}$$

Then, if 0 is special elliptic or special k -hyperbolic with k even, the tangent plane at 0 has the same orientation; if 0 is special elliptic or special k -hyperbolic with k odd the tangent space has opposite orientation.

2.8.7.

Proposition 9 (known for $n = 2$ [Bis65], here for $n \geq 3$). *Let S be a smooth, oriented, compact, 2-codimensional, real submanifold of \mathbb{C}^n whose all complex points are flat and special elliptic or special 1-hyperbolic. Then, on S , \sharp (special elliptic points) - \sharp (special 1-hyperbolic points) = $\chi(S)$. If S is a sphere, this number is 2.*

Proof. Let $p \in S$ be a complex point and π be the tangent hyperplane to S at p . Assume that

(**) *the orientation of S induces, on π , the orientation given by its complex structure,*
then $\pi \in H$.

If p is elliptic, the intersection number of H and $t(S)$ is 1; if p is 1-hyperbolic, the intersection number of H and $t(S)$ is -1 at p .

From the beginning of section 2.8.6, the sum of the intersection numbers of H and $t(S)$ at complex points p satisfying (**) is u_1 . Reversing the condition (**), and using Lemma 8, we get the Proposition. \square

3. PARTICULAR CASES: HORNED SPHERE; ELEMENTARY MODELS AND THEIR GLUING

3.1. We recall the following Harvey-Lawson theorem with real parameter to be used later.

3.1.1. Let $E \cong \mathbf{R} \times \mathbb{C}^{n-1}$, and $k : \mathbf{R} \times \mathbb{C}^{n-1} \rightarrow \mathbf{R}$ be the projection. Let $N \subset E$ be a compact, (oriented) CR subvariety of \mathbb{C}^{n+1} of real dimension $2n - 2$ and CR dimension $n - 2$, ($n \geq 3$), of class C^∞ , with negligible singularities (i.e. there exists a closed subset $\tau \subset N$ of $(2n - 2)$ -dimensional Hausdorff measure 0 such that $N \setminus \tau$ is a CR submanifold). Let τ' be the set of all points $z \in N$ such that either $z \in \tau$ or $z \in N \setminus \tau$ and N is not transversal to the complex hyperplane $k^{-1}(k(z))$ at z . Assume that N , as a current of integration, is d -closed and satisfies:

(H) there exists a closed subset $L \subset \mathbb{R}_{x_1}$ with $H^1(L) = 0$ such that for every $x \in k(N) \setminus L$, the fiber $k^{-1}(x) \cap N$ is connected and does not intersect τ' .

3.1.2.

Theorem 10 ([DTZ10] (see also [DTZ05])). *Let N satisfy (H) with L chosen accordingly. Then, there exists, in $E' = E \setminus k^{-1}(L)$, a unique C^∞ Levi-flat $(2n - 1)$ -subvariety M with negligible singularities in $E' \setminus N$, foliated by complex $(n - 1)$ -subvarieties, with the properties that M simply (or trivially) extends to E' as a $(2n - 1)$ -current (still denoted M) such that $dM = N$ in E' . The leaves are the sections by the hyperplanes $E_{x_1^0}$, $x_1^0 \in k(N) \setminus L$, and are the solutions of the “Harvey-Lawson problem” for finding a holomorphic subvariety in $E_{x_1^0} \cong \mathbb{C}^n$ with prescribed boundary $N \cap E_{x_1^0}$.*

3.1.3.

Remark 11. *Theorem 10 is valid in the space $E \cap \{\alpha_1 < x_1 < \alpha_2\}$, with the corresponding condition (H). Moreover, since N is compact, for convenient coordinate x_1 , we can assume $x_1 \in [0, 1]$.*

3.2. *To solve the boundary problem by Levi-flat hypersurfaces, S has to satisfy necessary and sufficient local conditions. A way to prove that these conditions can occur is to construct an example for which the solution is obvious.*

3.3. Sphere with one special 1-hyperbolic point (sphere with two horns): Example.

3.3.1. In \mathbb{C}^3 , let (z_j) , $j = 1, 2, 3$, be the complex coordinates and $z_j = x_j + iy_j$. In $\mathbb{R}^6 \cong \mathbb{C}^3$, consider the 4-dimensional subvariety (with negligible singularities) S defined by:

$$\begin{aligned} y_3 &= 0 \\ 0 \leq x_3 \leq 1; \quad &x_3(x_1^2 + y_1^2 + x_2^2 + y_2^2 + x_3^2 - 1) + (1 - x_3)(x_1^4 + y_1^4 + x_2^4 + y_2^4 + \\ &4x_1^2 - 2y_1^2 + x_2^2 + y_2^2) = 0 \\ -1 \leq x_3 \leq 0; \quad &x_3 = x_1^4 + y_1^4 + x_2^4 + y_2^4 + 4x_1^2 - 2y_1^2 + x_2^2 + y_2^2 \end{aligned}$$

The singular set of S is the 3-dimensional section $x_3 = 0$ along which the tangent space is not everywhere (uniquely) defined. S being in the real hyperplane $\{y_3 = 0\}$, the complex tangent spaces to S are $\{x_3 = x^0\}$ for convenient x^0 .

3.3.2. The tangent space to the hypersurface $f(x_1, y_1, x_2, y_2, x_3) = 0$ in \mathbb{R}^5 is

$$X_1 f'_{x_1} + Y_1 f'_{y_1} + X_2 f'_{x_2} + Y_2 f'_{y_2} + X_3 f'_{x_3} = 0,$$

Then, the tangent space to S in the hyperplane $\{y_3 = 0\}$ is:

for $0 \leq x_3$,

$$\begin{aligned} &2x_1[x_3 + 2(1 - x_3)(x_1^2 + 2)]X_1 + 2y_1[x_3 + 2(1 - x_3)(y_1^2 - 1)]Y_1 \\ &+ 2x_2[x_3 + (1 - x_3)(2x_2^2 + 1)]X_2 + 2y_2[x_3 + (1 - x_3)(2y_2^2 + 1)]Y_2 \\ &+ [(x_1^2 + y_1^2 + x_2^2 + y_2^2 + 3x_3^2 - 1) \\ &\quad - (x_1^4 + y_1^4 + x_2^4 + y_2^4 + 4x_1^2 - 2y_1^2 + x_2^2 + y_2^2)]X_3 = 0; \end{aligned}$$

for $x_3 \leq 0$,

$$4(x_1^2 + 2)x_1X_1 + 4(y_1^2 - 1)y_1Y_1 + 2(2x_2^2 + 1)x_2X_2 + 2(2y_2^2 + 1)y_2Y_2 - X_3 = 0.$$

3.3.3. The complex points of S are defined by the vanishing of the coefficients of X_j , $j=1,2,3,4$ in the equation of the tangent spaces

for $0 \leq x_3 \leq 1$,

$$\begin{aligned} x_1[x_3 + 2(1 - x_3)(x_1^2 + 2)] &= 0, \\ y_1[x_3 + 2(1 - x_3)(y_1^2 - 1)] &= 0, \\ x_2[x_3 + (1 - x_3)(2x_2^2 + 1)] &= 0, \\ y_2[x_3 + (1 - x_3)(2y_2^2 + 1)] &= 0. \end{aligned}$$

We have the solutions

h : $x_j = 0, y_j = 0, (j = 1, 2), x_3 = 0$;
 e_3 : $x_j = 0, y_j = 0, (j = 1, 2), x_3 = 1$.
 for $x_3 \leq 0$,

$$\begin{aligned}(x_1^2 + 2)x_1 &= 0, \\ (y_1^2 - 1)y_1 &= 0, \\ (2x_2^2 + 1)x_2 &= 0, \\ (2y_2^2 + 1)y_2 &= 0.\end{aligned}$$

We have the solutions

h : $x_j = 0, y_j = 0, (j = 1, 2), x_3 = 0$;
 e_1, e_2 : $x_1 = 0, y_1 = \pm 1, x_2 = 0, y_2 = 0, x_3 = -1$.

Remark that the tangent space to S at h is well defined. Moreover, the set S will be smoothed along its section by the hyperplane $\{x_3 = 0\}$ by a small deformation leaving h unchanged. In the following S will denote this smooth submanifold.

3.3.4.

Lemma 12. *The points e_1, e_2, e_3 are special elliptic; the point h is special $\{1\}$ -hyperbolic.*

Proof. Point e_3 : Let $x'_3 = 1 - x_3$, then the equation of S in the neighborhood of e_3 is:

$$(1 - x'_3)(x_1^2 + y_1^2 + x_2^2 + y_2^2 + x_3'^2 - 2x'_3) - x'_3(x_1^4 + y_1^4 + x_2^4 + y_2^4 + 4x_1^2 - 2y_1^2 + x_2^2 + y_2^2) = 0, \text{ i.e.}$$

$$2x'_3 = x_1^2 + y_1^2 + x_2^2 + y_2^2 + O(|z|^3), \text{ or } w = z\bar{z} + O(|z|^3)$$

then e_3 is special elliptic.

Points e_1, e_2 : Let $y'_1 = y_1 \pm 1, x'_3 = x_3 + 1$, then the equation of S in the neighborhood of e_1, e_2 is:

$$\begin{aligned}x'_3 - 1 &= x_1^4 + (y'_1 \mp 1)^4 + x_2^4 + y_2^4 + 4x_1^2 - 2(y'_1 \mp 1)^2 + x_2^2 + y_2^2 \\ &= x_1^4 + y_1'^4 \mp 4y_1'^3 + 6y_1'^2 \mp 4y_1' + 1 + x_2^4 + y_2^4 + 4x_1^2 - 2(y'_1 \mp 1)^2 + x_2^2 + y_2^2,\end{aligned}$$

then

$$x'_3 = x_1^4 + y_1'^4 \mp 4y_1'^3 + 4y_1'^2 + x_2^4 + y_2^4 + 4x_1^2 + x_2^2 + y_2^2, \text{ i.e.}$$

$$x'_3 = 4x_1^2 + 4y_1'^2 + x_2^2 + y_2^2 + O(|z|^3), \text{ or } w = 4z_1\bar{z}_1 + z_2\bar{z}_2,$$

then e_1, e_2 are special elliptic.

Point h : The equation of S in the neighborhood of h is:

for $x_3 \geq 0$,

$$\begin{aligned}x_3(x_1^2 + y_1^2 + x_2^2 + y_2^2 + x_3^2 - 1) \\ + (1 - x_3)(x_1^4 + y_1^4 + x_2^4 + y_2^4 + 4x_1^2 - 2y_1^2 + x_2^2 + y_2^2) = 0\end{aligned}$$

for $x_3 \leq 0$,

$$x_3 = x_1^4 + y_1^4 + x_2^4 + y_2^4 + 4x_1^2 - 2y_1^2 + x_2^2 + y_2^2, \text{ i.e.}$$

$x_3 = 4x_1^2 - 2y_1^2 + x_2^2 + y_2^2 + O(|z|^3)$, in both cases, up to the third order terms, i.e.: $w = z_1\bar{z}_1 + z_2\bar{z}_2 + 3\mathcal{R}e \ z_1^2$,

then h is special $\{1\}$ -hyperbolic. \square

3.3.5. *Section $\Sigma' = S \cap \{x_3 = 0\}$.* Up to a small smooth deformation, its equation is:

$$x_1^4 + y_1^4 + x_2^4 + y_2^4 + 4x_1^2 - 2y_1^2 + x_2^2 + y_2^2 = 0, \text{ in } \{x_3 = 0\}.$$

The tangent cone to Σ' at 0 is: $4x_1^2 - 2y_1^2 + x_2^2 + y_2^2 = 0$.

Locally, the section of S by the coordinate 3-space

$$x_1, y_1, x_3 \text{ is: } x_3 = 4x_1^2 - 2y_1^2 + O(|z|^3)$$

$$x_2, y_2, x_3 \text{ is: } x_3 = x_2^2 + y_2^2 + O(|z|^3)$$

3.3.1'. *Shape of $\Sigma' = S \cap \{x_3 = 0\}$ in the neighborhood of the origin 0 of \mathbb{C}^3 .*

Lemma 13. *Under the above hypotheses and notations,*

(i) $\Sigma = \Sigma' \setminus 0$ has two connected components σ_1, σ_2 .

(ii) *The closures of the three connected components of $S \setminus \Sigma'$ are submanifolds with boundaries and corners.*

Proof. (i) The only singular point of Σ' is 0. We work in the ball $B(0, A)$ of \mathbb{C}^2 (x_1, y_1, x_2, y_2) for small A and in the 3-space $\pi_\lambda = \{y_2 = \lambda x_2\}$, $\lambda \in \mathbb{R}$. For λ fixed, $\pi_\lambda \cong \mathbb{R}^3(x_1, y_1, x_2)$, and $\Sigma' \cap \pi_\lambda$ is the cone of equation $4x_1^2 - 2y_1^2 + (1 + \lambda^2)x_2^2 + O(|z|^3) = 0$ with vertex 0 and basis in the plane $x_2 = x_2^0$ the hyperboloid H_λ of equation $4x_1^2 - 2y_1^2 + (1 + \lambda^2)x_2^2 + O(|z|^3) = 0$; the curves H_λ have no common point outside 0. So, when λ varies, the surfaces $\Sigma' \cap \pi_\lambda$ are disjoint outside 0. The set Σ' is clearly connected; $\Sigma' \cap \{y_1 = 0\} = \{0\}$, the origin of \mathbb{C}^3 ; from above: $\sigma_1 = \Sigma \cap \{y_1 > 0\}$; $\sigma_2 = \Sigma \cap \{y_1 < 0\}$.

(ii) The three connected components of $S \setminus \Sigma'$ are the components which contain, respectively e_1, e_2, e_3 and whose boundaries are $\bar{\sigma}_1, \bar{\sigma}_2, \bar{\sigma}_1 \cup \bar{\sigma}_2$; these boundaries have corners as shown in the first part of the proof. \square

The connected component of $\mathbb{C}^2 \times \mathbb{R} \setminus S$ containing the point $(0, 0, 0, 0, 1/2)$ is the Levi-flat solution, the complex leaves being the sections by the hyperplanes $x_3 = x_3^0$, $-1 < x_3^0 < 1$.

The sections by the hyperplanes $x_3 = x_3^0$ are diffeomorphic to a 3-sphere for $0 < x_3^0 < 1$ and to the union of two disjoint 3-spheres for $-1 < x_3^0 < 0$, as can be shown intersecting S by lines through the origin in the hyperplane $x_3 = x_3^0$; Σ' is homeomorphic to the union of two 3-spheres with a common point.

The connected component of $\mathbb{C}^2 \times \mathbb{R} \setminus S$ containing the point $(0, 0, 0, 0, 1/2)$ is the Levi-flat solution, the complex leaves being the sections by the hyperplanes $x_3 = x_3^0$, $-1 < x_3^0 < 1$.

The sections by the hyperplanes $x_3 = x_3^0$ are diffeomorphic to a 3-sphere for $0 < x_3^0 < 1$ and to the union of two disjoint 3-spheres for $-1 < x_3^0 < 0$, as can be shown intersecting S by lines through the origin in the hyperplane $x_3 = x_3^0$; Σ' is homeomorphic to the union of two 3-spheres with a common point.

3.4. Sphere with one special 1-hyperbolic point (sphere with two horns). The example of section 3.3 shows that the necessary conditions of

section 2 can be realised. Moreover, from Proposition 2.8.7, the hypothesis on the number of complex points is meaningful.

3.4.1.

Proposition 14. [cf [Dol08]/[Proposition 2.6.1]] *Let $S \subset \mathbb{C}^n$ be a compact connected real 2-codimensional manifold such that the following holds:*

- (i) *S is a topological sphere; S is nonminimal at every CR point;*
- (ii) *every complex point of S is flat; there exist three special elliptic points $e_j, j = 1, 2, 3$ and one special 1-hyperbolic point h ;*
- (iii) *S does not contain complex manifolds of dimension $(n - 2)$;*
- (iv) *the singular CR orbit Σ' through h on S is compact and $\Sigma' \setminus \{h\}$ has two connected components σ_1 and σ_2 whose closures are homeomorphic to spheres of dimension $2n - 3$;*
- (v) *the closures S_1, S_2, S_3 of the three connected components S'_1, S'_2, S'_3 of $S \setminus \Sigma'$ are submanifolds with (singular) boundary.*

Then each $S_j \setminus \{e_j \cup \Sigma'\}$, $j = 1, 2, 3$ carries a foliation \mathcal{F}_j of class C^∞ with 1-codimensional CR orbits as compact leaves.

Proof. From conditions (i) and (ii), S satisfying the hypotheses of Proposition 1, near any elliptic flat point e_j , and of Proposition 6 near Σ' , all CR orbits being diffeomorphic to the sphere \mathbf{S}^{2n-3} . The assumption (iii) guarantees that all CR orbits in S must be of real dimension $2n - 3$. Hence, by removing small connected open saturated neighborhoods of all special elliptic points, and of Σ' , we obtain, from $S \setminus \Sigma'$, three compact manifolds S_j'' , $j = 1, 2, 3$, with boundary and with the foliation \mathcal{F}_j of codimension 1 given by its CR orbits whose first cohomology group with values in \mathbf{R} is 0, near e_j . It is easy to show that this foliation is transversely oriented. \square

3.4.2. Recall the Thurston's Stability Theorem ([CaC], Theorem 6.2.1).

Proposition 15. *Let (M, \mathcal{F}) be a compact, connected, transversely-orientable, foliated manifold with boundary or corners, of codimension 1, of class C^1 . If there is a compact leaf L with $H^1(L, \mathbf{R}) = 0$, then every leaf is homeomorphic to L and M is homeomorphic to $L \times [0, 1]$, foliated as a product,*

Then, from the above theorem, S_j'' is homeomorphic to $\mathbf{S}^{2n-3} \times [0, 1]$ with CR orbits being of the form $\mathbf{S}^{2n-3} \times \{x\}$ for $x \in [0, 1]$. Then the full manifold S_j is homeomorphic to a half-sphere supported by \mathbf{S}^{2n-2} and \mathcal{F}_j extends to S_j ; S_3 having its boundary pinched at the point h .

\square

3.4.3.

Theorem 16. *Let $S \subset \mathbb{C}^n$, $n \geq 3$, be a compact connected smooth real 2-codimensional submanifold satisfying the conditions (i) to (v) of Proposition 15. Then there exists a Levi-flat $(2n - 1)$ -subvariety $\tilde{M} \subset \mathbb{C} \times \mathbb{C}^n$ with boundary \tilde{S} (in the sense of currents) such that the natural projection $\pi :$*

$\mathbb{C} \times \mathbb{C}^n \rightarrow \mathbb{C}^n$ restricts to a bijection which is a CR diffeomorphism between \tilde{S} and S outside the complex points of S .

Proof. By Proposition 1, for every e_j , a continuous function ν'_j , C^∞ outside e_j , can be constructed in a neighborhood U_j of e_j , $j = 1, 2, 3$, and by Proposition 6, we have an analogous result in a neighborhood of Σ' . Furthermore, from Proposition 15, a smooth function ν''_j whose level sets are the leaves of \mathcal{F}_j can be obtained globally on $S'_j \setminus \{e_j \cup \Sigma'\}$. With the functions ν'_j and ν''_j , and analogous functions near Σ' , then using a partition of unity, we obtain a global smooth function $\nu_j: S_j \rightarrow \mathbf{R}$ without critical points away from the complex points e_j and from Σ' .

Let σ_1 , resp. σ_2 be the two connected, relatively compact components of $\Sigma \setminus \{h\}$, according to condition (iv); $\bar{\sigma}_1$, resp. $\bar{\sigma}_2$ are the boundary of S_1 , resp. S_2 , and $\bar{\sigma}_1 \cup \bar{\sigma}_2$ the boundary of S_3 . We can assume that the three functions ν_j are finite valued and get the same values on $\bar{\sigma}_1$ and $\bar{\sigma}_2$. Hence a function $\nu: S \rightarrow \mathbf{R}$.

The submanifold S being, locally, a boundary of a Levi-flat hypersurface, is orientable. We now set $\tilde{S} = N = \text{gr } \nu = \{(\nu(z), z) : z \in S\}$. Let $S_s = \{e_1, e_2, e_3, \bar{\sigma}_1 \cup \bar{\sigma}_2\}$.

$\lambda: S \rightarrow \tilde{S}$ ($z \mapsto \nu((z), z)$) is bicontinuous; $\lambda|_{S \setminus S_s}$ is a diffeomorphism; moreover λ is a CR map. Choose an orientation on S . Then N is an (oriented) CR subvariety with the negligible set of singularities $\tau = \lambda(S_s)$.

At every point of $S \setminus S_s$, $d_{x_1}\nu \neq 0$, then condition (H) (section 3.1.1) is satisfied at every point of $N \setminus \tau$.

Then all the assumptions of Theorem 10 being satisfied by $N = \tilde{S}$, in a particular case, we conclude that N is the boundary of a Levi-flat $(2n-2)$ -variety (with negligible singularities) \tilde{M} in $\mathbf{R} \times \mathbb{C}^n$.

Taking $\pi: \mathbb{C} \times \mathbb{C}^n \rightarrow \mathbb{C}^n$ to be the standard projection, we obtain the conclusion. \square

3.5. Generalizations: elementary models and their gluing.

3.5.1. The examples and the proofs of the theorems when S is homeomorphic to a sphere (sections 3.4) suggest the following definitions.

3.5.2. Definitions. Let T' be a smooth, locally closed (i.e. closed in an open set), connected submanifold of \mathbb{C}^n , $n \geq 3$. We assume that T' has the following properties:

- (i) T' is relatively compact, non necessarily compact, and of codimension 2.
- (ii) T' is nonminimal at every CR point.
- (iii) T' does not contain complex manifold of dimension $(n-2)$.
- (iv) T' has exactly 2 complex points which are flat and either special elliptic or special 1-hyperbolic.

(v) If $p \in T'$ is special 1-hyperbolic, the singular orbit Σ' through p is compact, $\Sigma' \setminus p$ has two connected components σ_1, σ_2 , whose closures are homeomorphic to spheres of dimension $2n - 3$.

(vi) If $p \in T'$ is special 1-hyperbolic, in the neighborhood of p , with convenient coordinates, the equation of T' , up to third order terms is

$$z_n = \sum_{j=1}^{n-1} (z_j \bar{z}_j + \lambda_j \operatorname{Re} z_j^2); \quad \lambda_1 > 1; \quad 0 \leq \lambda_j < 1 \quad \text{for } j \neq 1$$

or in real coordinates x_j, y_j with $z_j = x_j + iy_j$,

$$x_n = ((\lambda_1 + 1)x_1^2 - (\lambda_1 - 1)y_1^2) + \sum_{j=2}^{n-1} ((1 + \lambda_j)x_j^2 + (1 - \lambda_j)y_j^2) + O(|z|^3)$$

(vii) the closures, in T' , T_1, T_2, T_3 of the three connected components T'_1, T'_2, T'_3 of $T' \setminus \Sigma'$ are submanifolds with (singular) boundary. Let T''_j , $j = 1, 2, 3$ be neighborhoods of the T'_j in T' .

up- and down- 1-hyperbolic points. Let τ be the $(2n - 2)$ -submanifold with (singular) boundary contained into T' such that either $\bar{\sigma}_1$ (resp. $\bar{\sigma}_2$) is the boundary of τ near p , or Σ' is the boundary of τ near p . In the first case, we say that p is 1-up, (resp. 2-up), in the second that p is down. If T' is contained in a small enough neighborhood of Σ' in \mathbb{C}^n , such a T' will be called a *local elementary model*, more precisely it defines a *germ of elementary model around Σ* .

The union T of T_1, T_2, T_3 and of the germ of elementary model around the singular orbit at every special 1-hyperbolic point is called an *elementary model*. T behaves as a locally closed submanifold still denoted T .

3.5.3. Examples of elementary models. We will say that T is a *elementary model of type*:

- (a) if it has: two elliptic points;
- (b) if it has: one special elliptic point and one down- $\{1\}$ -hyperbolic point;
- (c₁) if it has: one special elliptic point and one 1-up- $\{1\}$ -hyperbolic point;
- (c₂) if it has: one special elliptic point and one 2-up- $\{1\}$ -hyperbolic point;
- (d₁) if it has: two special 1-up- $\{1\}$ -hyperbolic points;
- (d₂) if it has: two special 2-up- $\{1\}$ -hyperbolic points;
- (e) if it has: two special down- $\{1\}$ -hyperbolic points;

Other configurations are easily imagined.

The prescribed boundary of a Levi-flat hypersurface of \mathbb{C}^n in [DTZ05] and [DTZ10], whose complex points are flat and elliptic, is an elementary model of type (a).

3.5.4. Properties of elementary models. For instance, T is 1-up and has one special elliptic point, we solve the boundary problem as in S_1 in the proof of Theorem 16.

Proposition 17. *Let T be a local elementary model. Then, T carries a foliation \mathcal{F} of class C^∞ with 1-codimensional CR orbits as compact leaves.*

Proof. From the definition at the end of section 3.5.2 and Proposition 6. \square

3.5.5.

Theorem 18. *Let T be the elementary model there exists an open neighborhood T'' in T' carrying a smooth function $\nu : T'' \rightarrow \mathbb{R}$ whose level sets are the leaves of a smooth foliation.*

Proof. By removing small connected open saturated neighborhoods of every special elliptic point, and of Σ' , the singular orbit through every special 1-hyperbolic point p , we obtain, from $S \setminus \Sigma'$, three compact manifolds S_j'' , $j = 1, 2, 3$, with boundary,

(a) S_1 and S_2 containing one special elliptic point e or one special 1-hyperbolic point with the foliations $\mathcal{F}_1, \mathcal{F}_2$, from Propositions 1 and 17,

(b) S_3'' with the foliation \mathcal{F}_3 of codimension 1 given by its CR orbits whose first cohomology group with values in \mathbf{R} is 0, near e , or p . It is easy to show that this later foliation is transversely oriented.

From the Thurston's Stability Theorem (see section 3.4.2), S_3'' is homeomorphic to $\mathbf{S}^{2n-3} \times [0, 1]$, foliated as a product, with CR orbits being of the form $\mathbf{S}^{2n-3} \times \{x\}$ for $x \in [0, 1]$; hence smooth functions ν_1, ν_2, ν_3 , whose level sets are the leaves of the foliations $\mathcal{F}_1, \mathcal{F}_2, \mathcal{F}_3$ respectively, and using a partition of unity the desired function ν on T .

\square

3.6.

Theorem 19. *Let T be an elementary model. Then there exists a Levi-flat $(2n-1)$ -subvariety $\tilde{M} \subset \mathbb{C} \times \mathbb{C}^n$ with boundary \tilde{T} (in the sense of currents) such that the natural projection $\pi : \mathbb{C} \times \mathbb{C}^n \rightarrow \mathbb{C}^n$ restricts to a bijection which is a CR diffeomorphism between \tilde{T} and T outside the complex points of T .*

Proof. The submanifold T being, locally, a boundary of a Levi-flat hypersurface, is orientable. We now set $\tilde{T} = N = \text{gr } \nu = \{(\nu(z), z) : z \in S\} \subset E \cong \mathbf{R} \times \mathbb{C}^{n-1}$. Let T_s be the union of the flat complex points of T .

$\lambda : T \rightarrow \tilde{T}$ ($z \mapsto \nu((z), z)$) is bicontinuous; $\lambda|_{T \setminus T_s}$ is a diffeomorphism; moreover λ is a CR map. Choose an orientation on T . Then N is an (oriented) CR subvariety with the negligible set of singularities $\tau = \lambda(T_s)$.

Using Remark 11, at every point of $T \setminus T_s$, $d_{x_1} \nu \neq 0$, we see that condition (H) (section 3.1.1) is satisfied at every point of $N \setminus \tau$.

Then all the assumptions of Theorem 10 being satisfied by $N = \tilde{T}$, in a particular case, we conclude that N is the boundary of a Levi-flat $(2n-2)$ -variety (with negligible singularities) \tilde{M} in $\mathbf{R} \times \mathbb{C}^n$.

Taking $\pi : \mathbb{C} \times \mathbb{C}^n \rightarrow \mathbb{C}^n$ to be the standard projection, we obtain the conclusion. \square

3.7. Gluing of elementary models.

3.7.1. The gluing happens between two compatible elementary models along boundaries, for instance down and 1-up. Remark that the gluing can only be made at special 1-hyperbolic points. More precisely, it can be defined as follows.

The assumed properties of the submanifold S in section 2 in \mathbb{C}^n have a meaning in any complex analytic manifold X of complex dimension $n \geq 3$, and are kept under any holomorphic isomorphism.

We will define a submanifold S' of X obtained by gluing of elementary models by induction on the number m of models. An elementary model T in X is the image of an elementary model T_0 in \mathbb{C}^n by an analytic isomorphism of a neighborhood of T_0 in \mathbb{C}^n into X .

3.7.2. Let S' be a closed smooth real submanifold of X of dimension $2n - 2$ which is non minimal at every CR point. Assume that S' is obtained by gluing of m elementary models.

a) S' has a finite number of flat complex points, some special elliptic and the others special 1-hyperbolic;

b) for every special 1-hyperbolic p' , there exists a CR-isomorphism h induced by a holomorphic isomorphism of the ambient space \mathbb{C}^n from a neighborhood of p in T' onto a neighborhood of p' in S' .

c) for every CR-orbit $\Sigma_{p'}$ whose closure contains a special 1-hyperbolic point p' , there exists a CR-isomorphism h induced by a holomorphic isomorphism of the ambient space \mathbb{C}^n from a neighborhood of $\Sigma_p = \Sigma'_p \setminus p$ in T' onto a neighborhood V of $\Sigma_{p'}$ in S' .

Every special 1-hyperbolic point of S' which belongs to only one elementary model in S' will be called *free*.

We will define the gluing of one more elementary model to S' .

3.7.3. *Gluing an elementary model T of type (d_1) to a free down-1-hyperbolic point of S' .* Let h_1 be a CR-isomorphism from a neighborhood V_1 of $\bar{\sigma}'_1$ induced by a holomorphic isomorphism of the ambient space \mathbb{C}^n onto a neighborhood of σ_1 in S' . Let k_1 be a CR-isomorphism from a neighborhood T''_1 of T'_1 into X such that $k_1|_{V_1} = h_1$.

3.7.4.

Theorem 20. *The compact manifold or the manifold with singular boundary S' , obtained by the gluing of a finite number of elementary models, is the boundary of a Levi-flat hypersurface of X in the sense of currents.*

Proof. From Theorem 19 and the definition of gluing. \square

3.8. Examples of gluing. Denoting the gluing of the two models of type (d_1) and (d_2) to a free down-1-hyperbolic point of S' by: $\rightarrow (d_1) - (d_2)$, and the converse by: $(d_1) - (d_2) \rightarrow$, and, also, analogous configurations in the same way, we get:

torus: $(b) \rightarrow (d_1) - (d_2) \rightarrow (b)$; the Euler-Poincaré characteristic of a torus is $\chi(\mathbf{T}^k) = 0$: 2 special elliptic and 2 special 1-hyperbolic points.

bitorus: $(b) \rightarrow (d_1) - (d_2) \rightarrow (e) \rightarrow (d_1) - (d_2) \rightarrow (b)$.

4. CASE OF GRAPHS

(see [DTZ09] for the case of elliptic points only, and dropping the property of the function solution to be Lipschitz).

4.1. We want to add the following hypothesis: S is embedded into the boundary of a strictly pseudoconvex domain of \mathbb{C}^n , $n \geq 3$, and more precisely, let (z, w) be the coordinates in $\mathbb{C}^{n-1} \times \mathbb{C}$, with $z = (z_1, \dots, z_{n-1})$, $w = u + iv = z_n$, let Ω be a strictly pseudoconvex domain of $\mathbb{C}^{n-1} \times \mathbb{R}_u$ (i.e. the second fundamental form of the boundary $b\Omega$ of Ω is everywhere positive definite); let S be the graph $gr(g)$ of a smooth function $g : b\Omega \rightarrow \mathbb{R}_v$. notice that $b\Omega \times \mathbb{R}_v$ contains S and is strictly pseudoconvex.

Assume that S is a *horned sphere* (section 3.4), *satisfying the hypotheses of Theorem 16*. Denote by p_j , $j = i, \dots, 4$ the complex points of S . Our aim is to prove

4.2.

Theorem 21. *Let S be the graph of a smooth function $g : b\Omega \rightarrow \mathbb{R}_v$. Let $Q = (q_1, \dots, q_4) \in b\Omega$ be the projections of the complex points $P = (p_1, \dots, p_4)$ of S , respectively. Then, there exists a continuous function $f : \overline{\Omega} \rightarrow \mathbb{R}_v$ which is smooth on $\overline{\Omega} \setminus Q$ and such that $f|_{b\Omega} = g$, and $M_0 = \text{graph}(f) \setminus S$ is a smooth Levi flat hypersurface of \mathbb{C}^n . Moreover, each complex leaf of M_0 is the graph of a holomorphic function $\phi : \Omega' \rightarrow \mathbb{C}$ where $\Omega' \subset \mathbb{C}^{n-1}$ is a domain with smooth boundary (that depends on the leaf) and ϕ is smooth on $\overline{\Omega}'$.*

The natural candidate to be the graph M of f is $\pi(\tilde{M})$ where \tilde{M} and π are as in Theorem 16. We prove that this is the case proceeding in several steps.

4.3. Behaviour near S .

4.3.1. *Assume that D is a strictly pseudoconvex domain and that $S \subset bD$.*

Recall ([HL75][Theorem 10.4]: *Let D be a strictly pseudoconvex domain of \mathbb{C}^n , $n \geq 3$ with boundary bD , $\Sigma \subset bD$ be a compact connected maximally complex smooth $(2d - 1)$ -submanifold with $d \geq 2$. Then, Σ is the boundary of a uniquely determined relatively compact subset $V \subset \overline{D}$ such that $\overline{V} \setminus \Sigma$ is a complex analytic subset of D with finitely many singularities of pure*

dimension $\leq d - 1$, and near Σ , \bar{V} is a d -dimensional complex manifold with boundary.

V is said to be the solution of the boundary problem for Σ .

4.3.2.

Lemma 22 ([DTZ09]). *Let Σ_1, Σ_2 be compact connected maximally complex $(2d-1)$ -submanifolds of bD . Let V_1, V_2 be the corresponding solutions of the boundary problem. If $d \geq 2$, $2d \geq n + 1$ and $\Sigma_1 \cap \Sigma_2 = \emptyset$, then $V_1 \cap V_2 = \emptyset$.*

Let Σ be a CR orbit of the foliation of $S \setminus P$. Then Σ is a compact maximally complex $(2n-3)$ -dimensional real submanifold of \mathbb{C}^n contained in bD . Let $V = V_\Sigma$ be the solution of the boundary problem corresponding to Σ . From Theorem 16, $V = \pi(\tilde{V})$, where $\tilde{V} = (\tilde{M} \setminus \tilde{S}) \cap (\mathbb{C}^n \times \{x\})$ for suitable $x \in (0, 1)$, the projection on the x -axis being finite, we can always assume that it lies into $(0, 1)$. Moreover $\pi|_{\tilde{V}}$ is a biholomorphism $\tilde{V} \cong V$ and $M \setminus S \subset D$.

Let Σ_1, Σ_2 be two distinct orbits of the foliation of $S \setminus P$, and \bar{V}_1, \bar{V}_2 the corresponding leaves, then, from Lemma 22, $\bar{V}_1 \cap \bar{V}_2 = \emptyset$.

4.3.3. *Assume that S satisfies the full hypotheses of Theorem 21.*

Set $m_1 = \min_S g$, $m_2 = \max_S g$ and $r \gg 0$ such that

$$D = \Omega \times [m_1, m_2] \subset \subset \mathbf{B}(\mathbf{r}) \cap (\Omega \times i\mathbb{R}_v)$$

where $\mathbf{B}(\mathbf{r})$ is the ball $\{|(z, w)| < r\}$.

4.3.4.

Lemma 23. *Let $p \in S$ be a CR point. Then, near p , M is the graph of a function ϕ on a domain $U \subset \mathbb{C}_z^{n-1} \times \mathbb{R}_u$ which is smooth up to the boundary of U .*

Proof. Near p , each CR orbit Σ is smooth and can be represented as the graph of a CR function over a strictly pseudoconvex hypersurface and V_Σ as the graph of the local holomorphic extension of this function. From Hopf lemma, V is transversal to the strictly pseudoconvex hypersurface $d\Omega \times i\mathbb{R}_v$ near p . Hence the family of the V_Σ , near p , forms a smooth real hypersurface with boundary on S that is the graph of a smooth function ϕ from a relative open neighborhood U of p on $\bar{\Omega}$ into \mathbb{R}_v . Finally, Lemma 22 guarantees that this family does not intersect any other leaf V from M . \square

4.3.5.

Corollary 24. *If $p \in S$ is a CR point, each complex leaf V of M , near p , is the graph of a holomorphic function on a domain $\Omega_V \subset \mathbb{C}_z^{n-1}$, which is smooth up to the boundary of Ω_V .*

4.4. Solution as a graph of a continuous function.

4.4.1. Recall results of Shcherbina [Shc93] from:

(a) the Main Theorem:

Let G be a bounded strictly convex domain in $\mathbb{C}_z \times \mathbb{R}_u$ ($z \in \mathbb{C}$) and $\varphi : bG \rightarrow \mathbb{R}_v$ be a continuous function. Then the following properties hold, where $\Gamma = gr$, and $\hat{\Gamma}(\varphi)$ means polynomial hull of $\Gamma(\varphi)$:

(a_i) the set $\hat{\Gamma}(\varphi) \setminus \Gamma(\varphi)$ is the union of a disjoint family of complex discs $\{D_\alpha\}$;

(a_{ii}) for each α , there is a simply connected domain $\Omega_\alpha \subset \mathbb{C}_z$ and a holomorphic function $w = f_\alpha$, defined on Ω_α , such that D_α is the graph of f_α .

(a_{iii}) For each f_α , there exists an extension $f_\alpha^* \in C(\overline{\Omega}_\alpha)$ and $bD_\alpha = \{(z, w) \in b\Omega_\alpha \times \mathbb{C}_w : w = f_\alpha^*(z)\}$.

(b)

Lemma 25. Let $\{G_n\}_{n=0}^\infty$, $G_n \subset \mathbb{C}_z \times \mathbb{R}_u$, be a sequence of bounded strictly convex domains such that $G_n \rightarrow G_0$. Let $\{\varphi_n\}_{n=0}^\infty$, $\varphi_n : \partial G_n \rightarrow \mathbb{R}_v$ be a sequence of continuous functions such that $\Gamma(\varphi_n) \rightarrow \Gamma(\varphi_0)$ in the Hausdorff metric. Then, if Φ_n is the continuous function $\overline{G}_n \rightarrow \mathbb{R}_v$ such that $\hat{\Gamma}(\varphi) = \Gamma(\Phi)$, we have $\Gamma(\Phi_n) \rightarrow \Gamma(\Phi_0)$ in the Hausdorff metric.

(c)

Lemma 26. Let \mathcal{U} be a smooth connected surface which is properly embedded into some convex domain $G \subset \mathbb{C}_z \times \mathbb{R}_u$. Suppose that near each point of this surface, it can be defined locally by the equation $u = u(z)$. Then the surface \mathcal{U} can be represented globally as a graph of some function $u = U(z)$, defined on some domain $\Omega \subset \mathbb{C}_z$.

4.4.2.

Proposition 27. M is the graph of a continuous function $f : \overline{\Omega} \rightarrow \mathbb{R}_v$.

Proof. We will intersect the graph S with a convenient affine subspace of real dimension 4 to go back to the situation of Shcherbina.

Fix $a \in (\mathbb{C}_z^{n-1} \setminus 0)$ and, for a given point $(\zeta, \xi) \in \Omega$, with $\zeta \in \mathbb{C}_z^{n-1}$ and $\xi \in \mathbb{R}_u$, let $H_{(\zeta, \xi)} \subset \mathbb{C}_z^{n-1} \times \{\xi\}$ be the complex line through (ζ, ξ) in the direction $(a, 0)$. Set:

$$L_{(\zeta, \xi)} = H_{(\zeta, \xi)} + \mathbb{R}_u(0, 1), \quad \Omega_{(\zeta, \xi)} = L_{(\zeta, \xi)} \cap \Omega, \quad S_{(\zeta, \xi)} = (H_{(\zeta, \xi)} + \mathbb{C}_w(0, 1)) \cap S$$

Then $S_{(\zeta, \xi)}$ is contained in the strictly convex cylinder

$$(H_{(\zeta, \xi)} + \mathbb{C}_w(0, 1)) \cap (b\Omega \times i\mathbb{R}_v)$$

and is the graph of $g|_{b\Omega_{(\zeta, \xi)}}$.

From (a_{ii}), the polynomial hull of $S_{(\zeta, \xi)}$ is a continuous graph over $\overline{\Omega}_{(\zeta, \xi)}$. Consider $M = \pi(\tilde{M})$ and set

$$M_{\zeta, \xi} = (H_{(\zeta, \xi)} + \mathbb{C}_w(0, 1)) \cap M.$$

It follows that $M_{\zeta, \xi}$ is contained in the polynomial hull $\hat{S}_{(\zeta, \xi)}$. From (a_{iii}), $\hat{S}_{(\zeta, \xi)}$ is a graph over $\overline{\Omega}_{(\zeta, \xi)}$ foliated by analytic discs, so $M_{\zeta, \xi}$ is a graph over a subset U of $\overline{\Omega}_{(\zeta, \xi)}$.

Every analytic disc Δ of $\hat{S}_{(\zeta, \xi)}$ had its boundary on $S_{(\zeta, \xi)}$. Since all the complex points of S are isolated, $b\Delta$ contains a CR point p of S ; from Lemma 23, near p , $M_{\zeta, \xi}$ is a graph over $\overline{\Omega}_{(\zeta, \xi)}$. Near p , Δ is contained in $M_{\zeta, \xi}$, then in a closed complex analytic leaf V_Σ of M ; so $\Delta \subset V_\Sigma \subset M$; but $\Delta \subset H_{(\zeta, \xi)} + \mathbb{C}_w(0, 1)$; then: $\Delta \subset M_{\zeta, \xi}$. Consequently, near p , $M_{\zeta, \xi} = \hat{S}_{(\zeta, \xi)}$.

It follows that M is the graph of a function $f : \overline{\Omega} \rightarrow \mathbb{R}_v$.

One proves, using (b), that f is continuous on Ω , whence on $\overline{\Omega} \setminus Q$, by Lemma 23. Then continuity at every q_j is proved using the *Kontinuitätsatz* on the domain of holomorphy $\Omega \times i\mathbb{R}_v$. \square

4.5. Regularity. The property: $M \setminus P = (p_1, \dots, p_4)$ is a smooth manifold with boundary results from:

4.5.1.

Lemma 28. *Let U be a domain of $\mathbb{C}_z^{n-i} \times \mathbb{R}_u$, $n \geq 2$, $f : U \rightarrow \mathbb{R}_v$ a continuous function. Let $A \subset \text{graph}(f)$ be a germ of complex analytic set of codimension 1. Then A is a germ of complex manifold which is a graph of over \mathbb{C}_z^{n-i} .*

Proof. Assume that A is a germ at 0. Let $g \in \mathcal{O}$, $h \neq 0$ such that $A = \{h = 0\}$. For $\varepsilon \ll 1$, let \mathbf{D}_ε be the disc $\{z = 0\} \cap \{|w| < \varepsilon\}$, then $A \cap \mathbf{D}_\varepsilon = \{0\}$, i.e. A is w -regular.

Let $\pi : \mathbb{C}_{z,w}^n \rightarrow \mathbb{C}_z^{n-1}$ be the projection. The local structure theorem for analytic sets gives:

for some neighborhood U of 0 in \mathbb{C}_z^{n-1} , there exists an analytic hypersurface $\Delta \subset U$ such that: $A_\Delta = A \cup ((U \setminus \Delta) \times \mathbf{D}_\varepsilon)$ is a manifold;

$\pi/A_\Delta \rightarrow U \setminus \Delta$ is a $d(\in \mathbb{N})$ -sheeted covering.

It is easy to show that the covering $\pi : A_\Delta \rightarrow U \setminus \Delta$ is trivial.

Then we may define d holomorphic functions $\tau_1, \dots, \tau_d : U \setminus \Delta \rightarrow \mathbb{C}$ such that A_Δ is the union of the graphs of the τ_j . By the Riemann extension theorem, the functions τ_j extend as holomorphic functions $\tau_j \in \mathcal{O}(U)$. Suppose that $\tau_j \neq \tau_k$, for $j \neq k$, then for some disc $\mathbf{D} \subset U$ centered at 0, we have $\tau_j|_{\mathbf{D}} \neq \tau_k|_{\mathbf{D}}$, then $(\tau_j - \tau_k)|_{\mathbf{D}}$ vanishes only at 0. But, from the hypothesis, in restriction to \mathbf{D} , $\{Re(\tau_j - \tau_k) = 0\} \subset \{\tau_j - \tau_k = 0\}|_{\mathbf{D}} = \{0\}$, impossible. \square

4.6.

Proof of the Theorem 21. Consider the foliation of $S \setminus P$ given by the level sets of the smooth function $\nu : S \rightarrow [0, 1]$ (sections 2.3 and 2.7) and set $L_t = \{\nu = t\}$ for $t \in (0, 1)$. Let $V_t \subset \overline{\Omega} \times i\mathbb{R}_v \subset \mathbb{C}^n$ be the complex leaf of M bounded by L_t .

By Proposition 27, M is the graph of a continuous function over Ω , and, by Lemma 28, each leaf V_t is a complex smooth hypersurface and $\pi|_{V_t}$ is a submersion.

→

Since Ω is strictly convex, as in Shcherbina (see 4.4.1, c)), $\pi|_{V_t}$ is 1-1, then, by Corollary 24, π sends V_t onto a domain $\Omega_t \subset \mathbb{C}_z^{n-1}$ with smooth boundary. Let

$$\pi_u : (\mathbb{C}_z^{n-1} \times \mathbb{R}_u) \times i\mathbb{R}_v \rightarrow \mathbb{R}_u$$

$$\pi_v : (\mathbb{C}_z^{n-1} \times \mathbb{R}_u) \times i\mathbb{R}_v \rightarrow \mathbb{R}_v$$

then $\pi_u|_{L_t} = a_t \cdot \pi|_{L_t}$ and $\pi_v|_{L_t} = b_t \cdot \pi|_{L_t}$ where a_t, b_t are smooth functions on $b\Omega_t$. Moreover $b\Omega_t, a_t, b_t$ depend smoothly on t .

If $(z_t, w_t) \in M$, then w_t varies on V_t , so w_t is the holomorphic extension of $a_t + ib_t$ to Ω_t . In particular u_t and v_t are smooth in (z, t) , from the Bochner-Martinelli formula.

$\frac{\partial u_t}{\partial t}$ is harmonic on Ω_t for each t and has a smooth extension on $b\Omega_t$.

From Lemma 23 and Corollary 24, $\frac{\partial u_t}{\partial t}$ does not vanish on $b\Omega_t$. Since the CR orbits L_t are connected from Proposition 14, $b\Omega_t$ is also connected, hence $\frac{\partial u_t}{\partial t}$ has constant sign on $b\Omega_t$. Then, by the maximum principle, also $\frac{\partial u_t}{\partial t}$ on Ω_t and, in particular does not vanish. This implies that $M \setminus S$ is the graph of a smooth function over Ω which smoothly extends to $\overline{\Omega} \setminus Q$.

From Proposition 27, M is the graph of a continuous function over $\overline{\Omega}$. \square

4.7. Elementary smooth models.

4.7.1. Definition. An elementary smooth model in \mathbb{C}^n is an elementary model in the sense of section 3.5.2 and satisfying the further condition which makes sense from Theorem 21:

(G) Let (z, w) be the coordinates in $\mathbb{C}^{n-1} \times \mathbb{C}$, with $z = (z_1, \dots, z_{n-1}), w = u + iv = z_n$, let Ω be a strictly pseudoconvex domain of $\mathbb{C}^{n-1} \times \mathbb{R}_u$; assume that T' is the graph of a smooth function $g : b\Omega \rightarrow \mathbb{R}_v$.

4.7.2.

Theorem 29. Let T be an elementary smooth model. Then, there exists a continuous function $f : \overline{\Omega} \rightarrow \mathbb{R}_v$ which is smooth on $\overline{\Omega} \setminus Q$ and such that $f|_{b\Omega} = g$, and $M_0 = \text{graph}(f) \setminus S$ is a smooth Levi flat hypersurface of \mathbb{C}^n ; in particular, S is the boundary of the hypersurface $M = \text{graph}(f)$

Proof. similar to the proof of Theorem 21. \square

4.7.3. Gluing of elementary smooth models. In an open set of \mathbb{C}^n , a coordinate system (z, w) of $\mathbb{C}_z^{n-1} \times \mathbb{R}_u$ defines an $(n-1, 1)$ -frame.

To define the gluing of elementary models (section 3.7) we considered a CR-isomorphism from an open set of \mathbb{C}^n induced by a holomorphic isomorphism of the ambient space \mathbb{C}^n onto a an open set of \mathbb{C}^n . To define the

gluing of elementary smooth models, we have to consider a holomorphic isomorphism of the ambient space \mathbb{C}^n onto an open set of \mathbb{C}^n sending an $(n-1, 1)$ -frame of $\mathbb{C}_z^{n-1} \times \mathbb{R}_u$ onto an $(n-1, 1)$ -frame of $\mathbb{C}_{z'}^{n-1} \times \mathbb{R}_{u'}$.

As in section 3.7.1, we will define a submanifold S' of X obtained by gluing of elementary smooth models by induction on the number m of models. An elementary smooth model T in X is the image of an elementary smooth model T_0 of \mathbb{C}^n by an analytic isomorphism of a neighborhood of T_0 in \mathbb{C}^n into X .

Gluing an elementary smooth model T of type (d_1) to a free down-1-hyperbolic point of S' .

Every elementary smooth model is contained in a cylinder $b\Omega \times \mathbb{R}_v$ determined by Ω and an $(n-1, 1)$ -frame. Two sets Ω are *compatible* if either they coincide or one is part of the other.

The announced gluing is defined in the following way: there exists a CR-isomorphism h_1 from a neighborhood V_1 of $\bar{\sigma}'_1$ induced by a holomorphic isomorphism of the ambient space \mathbb{C}^n onto a neighborhood of σ_1 in S' . Let k_1 be a CR-isomorphism from a neighborhood T''_1 of T'_1 into X such that $k_1|_{V_1} = h_1$, and there exists a common $(n-1, 1)$ -frame on which the corresponding sets Ω are compatible. The existence of such a situation is possible as the example of the horned (almost everywhere) smooth sphere shows (Theorem 21.).

Remark that the gluing implies that the obtained submanifold S' is C^0 and smooth except at the complex points.

Other gluing are obtained in a similar way. Hence:

Theorem 30. *The manifold S' obtained by gluing of elementary smooth models is of class C^0 , and smooth except at the complex points.*

Corollary 31. *The manifold S' is the boundary of manifold M of class C^∞ whose interior is a Levi-flat smooth hypersurface.*

REFERENCES

- [Bis65] Errett Bishop, *Differentiable manifolds in complex Euclidean space*, Duke Math. J. **32** (1965), 1–21. MR 0200476 (34 #369)
- [BK91] Eric Bedford and Wilhelm Klingenberg, *On the envelope of holomorphy of a 2-sphere in \mathbb{C}^2* , J. Amer. Math. Soc. **4** (1991), no. 3, 623–646. MR 1094437 (92j:32034)
- [CS51] Shiing-shen Chern and E. Spanier, *A theorem on orientable surfaces in four-dimensional space*, Comment. Math. Helv. **25** (1951), 205–209. MR 0044883 (13,492d)
- [Dol08] Pierre Dolbeault, *On Levi-flat hypersurfaces with given boundary in \mathbb{C}^n* , Sci. China Ser. A **51** (2008), no. 4, 541–552. MR 2395404 (2009c:32067)
- [Dol11] P. Dolbeault, *Complex plateau problem: old and new results and prospects*, Arxiv preprint arXiv:1105.4417 (2011).
- [DTZ05] Pierre Dolbeault, Giuseppe Tomassini, and Dmitri Zaitsev, *On boundaries of Levi-flat hypersurfaces in \mathbb{C}^n* , C. R. Math. Acad. Sci. Paris **341** (2005), no. 6, 343–348. MR 2169149 (2006e:32048)

- [DTZ09] P. Dolbeault, G. Tomassini, and D. Zaitsev, *Boundary problem for Levi flat graphs*, Arxiv preprint arXiv:0912.1332 (2009).
- [DTZ10] Pierre Dolbeault, Giuseppe Tomassini, and Dmitri Zaitsev, *On Levi-flat hypersurfaces with prescribed boundary*, Pure Appl. Math. Q. **6** (2010), no. 3, Special Issue: In honor of Joseph J. Kohn. Part 1, 725–753. MR 2677311 (2011f:32074)
- [HL75] F. Reese Harvey and H. Blaine Lawson, Jr., *On boundaries of complex analytic varieties. I*, Ann. of Math. (2) **102** (1975), no. 2, 223–290. MR 0425173 (54 #13130)
- [Lai72] Hon Fei Lai, *Characteristic classes of real manifolds immersed in complex manifolds*, Trans. Amer. Math. Soc. **172** (1972), 1–33. MR 0314066 (47 #2618)
- [Lai74] ———, *On the topology of the even-dimensional complex quadrics*, Proc. Amer. Math. Soc. **46** (1974), 419–425. MR 0358846 (50 #11305)
- [Pol08] P. Polo, *Grassmanniennes orientées réelles*, Personnel communication, 1 fév. 2008.
- [Shc93] N. V. Shcherbina, *On the polynomial hull of a graph*, Indiana Univ. Math. J. **42** (1993), no. 2, 477–503. MR 1237056 (95e:32017)
- [Ste99] Norman Steenrod, *The topology of fibre bundles*, Princeton Landmarks in Mathematics, Princeton University Press, Princeton, NJ, 1999, Reprint of the 1957 edition, Princeton Paperbacks. MR 1688579 (2000a:55001)

INSTITUT DE MATHÉMATIQUES DE JUSSIEU, UPMC, 4, PLACE JUSSIEU 75005 PARIS
E-mail address: pierre.dolbeault@upmc.fr